

A class of quasi-stationary regular line elements for the Schwarzschild geometry

Kayll Lake
Department of Physics
Queen's University
Kingston Ontario Canada K7L 3N6
lake@astro.queensu.ca

3 July 1994

Abstract

A one-fold infinity of explicit quasi-stationary regular line elements for the Schwarzschild geometry is obtained directly from the vacuum Einstein equations. The class includes the familiar Eddington-Finkelstein coordinates, and the coordinates discussed recently by Kraus and Wilczek.

The quasi-stationary spherically symmetric line element in curvature coordinates (labelled (t, r, θ, ϕ)) is given by

$$ds^2 = -a(r)dt^2 + 2b(r)dtdr + c(r)dr^2 + r^2d\Omega^2 \quad (1)$$

where $d\Omega^2$ is the metric of a unit sphere ($d\theta^2 + \sin^2(\theta)d\phi^2$). The term “curvature coordinates” is derived from the coefficient of $d\Omega^2$, and the term “quasi-stationary” is derived from the fact that for the metric (1), $\xi^\alpha = \delta^\alpha_t$ is a Killing vector that is timelike for $a(r) > 0$.

The purpose of this note is to solve the vacuum Einstein equations

$$R^\alpha_\beta = 0 \quad (2)$$

in terms of the coordinates defining the line element (1). The system is over determined and any one of $a(r)$, $b(r)$ or $c(r)$ can be prescribed. We content ourselves here by simply specifying constants.

With $c(r) = c = \text{const.} \geq 0$, we solve (2) and obtain (in geometrical units)

$$ds^2 = -(1 - 2m/r)dt^2 + 2\epsilon(1 - c + 2mc/r)^{\frac{1}{2}}dtdr + cdr^2 + r^2d\Omega^2 \quad (3)$$

where $\epsilon = \pm 1$. The constant m is the effective gravitational mass. The constant c has no invariant physical significance. (Whereas the surfaces $t = \text{constant}$

have distinct properties for different choices of c , this is merely a choice of slicing of the spacetime.) With $0 \leq c \leq 1$, the r - t subspaces of the metric (3) are free of coordinate singularities over the range $0 < r < \infty$. The familiar Eddington-Finkelstein coordinates are obtained by setting $c = 0$. The coordinates discussed, and exploited in a modern context, recently by Kraus and Wilczek [1] are obtained by setting $c = 1$. These coordinates have an interesting history [2]. I am not aware of a previous discussion of the class given by (3). (However, in view of the elementary nature of the problem, and solution, it would be somewhat of a surprise if the form (3) is actually new!) Whereas with $0 \leq c \leq 1$ the line element (3) is free of coordinate singularities, the coordinates are incomplete. The completion can be carried out for $0 \leq c \leq 1$ exactly analogously to the case $c = 0$ that is thoroughly treated in standard texts.

The procedure that yielded the form (3) fails to give anything of further interest. In particular, setting $a(r) = 1$ (by choice of scale of t) leads, with (2), only to Minkowski space. The forms obtained with $b(r) = b = \text{const.} \geq 0$ all have the familiar coordinate singularity at $r = 2m$.

This work was supported by a grant from the National Sciences and Engineering Research Council of Canada. The entire calculation reported here was executed in a few CPU seconds on a notebook computer running GRTensor under MapleV Release 3. GRTensor runs on any hardware platform which supports Maple and is available free of charge from the author.

References

- [1] P. Kraus and F. Wilczek, "Some Applications of a Simple Line Element for the Schwarzschild Geometry", preprint gr-qc/9406042.
- [2] A sketch of the history is given by W. Israel "Dark Stars: the evolution of an idea" in *300 Years of Gravitation*, ed. by S. W. Hawking and W. Israel (Cambridge University Press, Cambridge, 1987). A detailed history of the early years is given by J. Eisenstaedt, "Histoire et Singularités de la Solution de Schwarzschild (1915-1923)" in *Archive for History of Exact Sciences*, Volume 27, Number 2, 1982.